ELSEVIER

Volumes 271–273, 12 June 1998

Journal of ALLOYS AND COMPOUNDS

Contents

Preface xv
Organisers and sponsors xvi

The future of nuclear energy in the world 1
B. Barré (Paris, France)

The bioinorganic chemistry of actinides in blood 6
D.M. Taylor (Karlsruhe, Germany and Cardiff, UK)

Actinide Detection, Monitoring, Analysis and Characterization

Strengthening IAEA safeguards through environmental sampling and analysis 11
D.L. Donohue (Vienna, Austria)

The use of SIMS for uranium localization in biological research 19
A. Amaral (Recife Pernambuco, Brazil), P. Galle, F. Escaig (Créteil, France), C. Cossonnet, M.H. Henge-Napoli, E. Ansoborlo (Fontenay-aux-Roses, France), L. Zhang (Créteil, France)

Alpha-spectrometry measurement of Am and Cm at trace levels in environmental samples using extraction chromatography . . 25
F. Goutelard, M. Morello, D. Calmet (Orsay, France)

Simple on-line monitoring of α- and β-emitters by solid scintillation counting 31
M. Lochny, W. Ullrich, U. Wenzel (Jülich, Germany)

Chemical composition of inhalable MOX powders prepared according to a SOLGEL process: energy dispersive X-ray analysis of individual particles 38
P. Massiot, B. Leprince, C. Lizon, G. Rateau (Bruyères-le-Châtel, France), K. Richter (Karlsruhe, Germany), P. Fritsch (Bruyères-le-Châtel, France)

A fast method for low-level actinide measurement in concrete 42
C. Nebelung, H. Nitsche, G. Bernhard (Dresden, Germany)

Trace analysis of plutonium in environmental samples by resonance ionization mass spectroscopy (RIMS) 45
M. Nunnemann, N. Erdmann, H.-U. Hasse, G. Huber, J.V. Kratz, P. Kunz, A. Mansel, G. Passler, O. Stetzer, N. Trautmann, A. Waldek (Mainz, Germany)

Separation of actinides from a bone ash matrix with extraction chromatography 49
R. Pilviö, M. Bickel (Geel, Belgium)

Exposure limits and assessment of intake for inhaled soluble uranium compounds 54
G.N. Stradling, J.C. Moody, A. Hodgson, T.P. Fell, E. Rance (Didcot, UK)

Simultaneous counting of radiation emitted from actinides with improved phoswich detectors by applying an optical filter . . . 58
S. Usuda, K. Yasuda, S. Sakurai (Ibaraki-ken, Japan)

Determination of actinide isotope ratios using glow discharge optogalvanic spectroscopy 62
J.P. Young, R.W. Shaw, C.M. Barshick, J.M. Ramsey (Oak Ridge, TN, USA)

Biochemistry of the Actinides

Issues and research on the biochemistry of inhaled actinides 66
R.A. Guilmette (Albuquerque, NM, USA)

Decorporation of actinides: a review of recent research 72
G.N. Stradling (Didcot, UK)

Biotransformation of uranium and other actinides in radioactive wastes 78
A.J. Francis (Upton, NY, USA)

The bioinorganic chemistry of Np, Pu and Am in mammalian liver 85
F. Paquet (Fontenay-aux-Roses, France), B. Ramounet (Bruyères-le-Châtel, France), H. Métivier (Fontenay-aux-Roses, France), D.M. Taylor (Cardiff, UK)

Elsevier Science S.A.

Experimental approaches to improve the available chelator treatments for Np decorporation 89
P. Fritsch, B. Ramounet (Bruyères-le-Châtel, France), R. Burgada (Paris, France), H. Métivier, F. Paquet (Pierrelatte, France), J.L. Poncy (Bruyères-le-Châtel, France)

The effect of a delayed chronic DTPA treatment on the Pu microdistribution in rat bone: preliminary results using hemi-blocks of femurs 93
H. Le Naour, B. Ramounet, L. Le Foll, P. Fritsch (Bruyères-le-Châtel, France)

In vitro measurement of UO_2/phosphate ion interactions at the particle surface: preliminary results using electron dispersive energy (EDS) and ^{32}P adsorption assays 96
S. Matton (Bruyères-le-Châtel, France), J.-P. Gallien (Gif-sur-Yvette, France), P. Massiot, I. Bailly, P. Fritsch (Bruyères-le-Châtel, France)

Extraction of Pu from rat bone by formic acid/guanidine treatment after intravenous ^{238}Pu-citrate administration: influence of time after contamination and of a chronic DTPA treatment 100
B. Ramounet, P. Fritsch (Bruyères-le-Châtel, France)

Early biokinetics of actinides after intravenous administration of soluble forms of Pu(IV), Np(IV) and Np(V) 103
B. Ramounet, S. Matton, G. Grillon, P. Fritsch (Bruyères-le-Châtel, France)

Investigations by time-resolved laser-induced fluorescence and capillary electrophoresis of the uranyl–phosphate species: application to blood serum 106
S. Scapolan, E. Ansoborlo (Fontenay-aux-Roses, France), C. Moulin (Gif-sur-Yvette, France), C. Madic (Bagnols-sur-Cèze, France)

Computer modelling of the chemical speciation of lanthanide and actinide elements in the human gastrointestinal tract 112
L.M. Webb, D.M. Taylor, D.R. Williams (Cardiff, UK)

Coordination Chemistry–Organometallics

Designing new catalytic C–C and C–N bond formations promoted by organoactinides 116
M.S. Eisen, T. Straub, A. Haskel (Haifa, Israel)

Determination of structural parameters of uranyl ions complexed with organic acids using EXAFS 123
M.A. Denecke, T. Reich, M. Bubner, S. Pompe, K.H. Heise, H. Nitsche (Dresden, Germany), P.G. Allen, J.J. Bucher, N.M. Edelstein, D.K. Shuh (Berkeley, CA, USA)

Electronic structures of actinyl nitrate–triethyl phosphate complexes using the DV–DS method 128
M. Hirata (Ibaraki, Japan), R. Sekine (Shizuoka, Japan), J. Onoe (Saitama, Japan), H. Nakamatsu, T. Mukoyama (Kyoto, Japan), K. Takeuchi (Saitama, Japan), S. Tachimori (Ibaraki, Japan)

Crown ether complexes of $UO_2(NCS)_2$ and $Th(NCS)_4$: clues to solution behavior or just interesting supramolecular structures? 133
R.D. Rogers, J. Zhang, D.T. Campbell (Tuscaloosa, AL, USA)

Structural studies on monomeric and dimeric uranyl bis(dimethylmalonato) complexes 139
Y. Zhang, D. Collison, F.R. Livens, M. Helliwell (Manchester, UK), H. Eccles, N. Tinker (Preston, UK)

Uranium amides as precursors to cationic and/or pentavalent compounds 144
M. Ephritikhine, J.C. Berthet, C. Boisson, M. Lance, M. Nierlich (Gif-sur-Yvette, France)

Triamidoamine complexes of the actinides; from mixed-valence bimetallics to dinitrogen activation 150
P. Roussel (Coventry, UK), N.D. Tinker (Preston, UK), P. Scott (Coventry, UK)

Peculiarities of complex formation of hexavalent Np and Pu with tetrahedral XO_4^{2-} oxoanions (X=S, Se, Cr, Mo) 154
A. Fedosseev, N. Budantseva, A. Bessonov, M. Grigoriev (Moscow, Russia), J.C. Krupa (Orsay, France)

Trivalent minor actinides/lanthanides separation, using organophosphinic acids 159
C. Hill, C. Madic, P. Baron (Bagnols-sur-Cèze, France), M. Ozawa, Y. Tanaka (Ibaraki-ken, Japan)

Electrochemical studies on bis(β-diketonato)(dimethyl sulfoxide)dioxouranium(VI) in dimethyl sulfoxide 163
K. Mizuguchi, S.-H. Lee (Tokyo, Japan), Y. Ikeda (Chiba, Japan), H. Tomiyasu (Tokyo, Japan)

Complex formation of divalent lanthanides and actinides 168
S.A. Kulyukhin (Moscow, Russia)

Coordination chemistry and extraction properties of phosphonopyridyl N,P oxides 172
E.M. Bond, X. Gan (Albuquerque, NM, USA), J.R. Fitzpatrick (Los Alamos, NM, USA), R.T. Paine (Albuquerque, NM, USA)

Coordination chemistry of neptunyl(V) compounds by means of Mössbauer spectroscopy 176
M. Saeki, M. Nakada, T. Nakamoto, N.M. Masaki, T. Yamashita (Ibaraki, Japan)

Uranyl monopicolinate complexes 180
P.R. Silverwood, D. Collison, F.R. Livens, R.L. Beddoes (Manchester, UK), R.J. Taylor (Sellafield, UK)

Structural study of holmium (III) and uranium (VI) organic ligand complexes by extended X-ray absorbtion fine-structure spectroscopy 184
T. Yaita, H. Narita, S. Suzuki, S. Tachimori (Ibaraki, Japan), H. Shiwaku, H. Motohashi (Hyogo, Japan)

Environmental Remediation, Waste Management and Actinide Mitigation

A method to predict free energies of formation of mineral phases in the U(VI)–SiO_2–H_2O system 189
S.B. Clark (Pullman, WA, USA), R.C. Ewing (Ann Arbor, MI, USA), J.C. Schaumloffel (Pullman, WA, USA)

Direct disposal of spent nuclear fuel: comparison between experimental and modelled actinide solubiluties in natural waters . . 194
L.O. Werme, K. Spahiu (Stockholm, Sweden)

Uranium speciation in waters of different uranium mining areas 201
G. Bernhard, G. Geipel, V. Brendler, H. Nitsche (Dresden, Germany)

Complexation of Th (IV) with the siderophore pyoverdine A 206
M. Bouby, I. Billard, J. MacCordick (Strasbourg, France)

Oxidation kinetics of plutonium in air: consequences for environmental dispersal 211
J.M. Haschke, T.H. Allen, J.C. Martz (Los Alamos, NM, USA)

Up-to-date radioecological situation around the 'Mayak' nuclear facility 216
B.F. Myasoedov, E.G. Drozhko (Moscow, Russia)

The incorporation of plutonium and neptunium in zirconolite and perovskite 221
B.D. Begg, E.R. Vance (Menai, Australia), S.D. Conradson (Los Alamos, NM, USA)

Lysimeter studies to investigate the leaching of ^{241}Am from low level radioactive waste 227
E.J. Kelly, D.I. Bell (Sellafield, U.K), T.L. Johnstone (Risley, U.K)

Capture of Pu(V), Np(V) and Pu(VI) from alkaline solutions by hydroxides of Pu(IV), Th(IV) and La(III) 231
N. Budantseva, A. Bessonov, I. Tananaev, A. Fedosseev (Moscow, Russia), C. Delegard (Richland, WA, USA)

Actinides immobilization in new matrices based on solid solutions: $Th_{4-x}M^{IV}_x(PO_4)_4P_2O_7$, ($M^{IV}$=$^{238}U$,$^{239}Pu$) 236
N. Dacheux, R. Podor, B. Chassigneux, V. Brandel, M. Genet (Orsay, France)

U and Pu L_{iii} XAFS of Pu-doped glass and ceramic waste forms 240
N.J. Hess, W.J. Weber (Richland, WA, USA), S.D. Conradson (Los Alamos, NM, USA)

Applicability of insoluble tannin to treatment of waste containing americium 244
T. Matsumura, S. Usuda (Ibaraki-ken, Japan)

The separation of trivalent actinides from lanthanides by dithiophosphinic acids from HNO_3 acid medium 248
G. Modolo, R. Odoj (Jülich, Germany)

Surface sorption and surface diffusion of NpO_2^+ with poorly crystallized ferric oxide 252
S. Nagasaki, S. Tanaka, M. Todoriki, A. Suzuki (Tokyo, Japan)

Actinide immobilization in the subsurface environment by in-situ treatment with a hydrolytically unstable organophosphorus complexant: Uranyl uptake by calcium phytate 257
K.L. Nash, M.P. Jensen, M.A. Schmidt (Argonne, IL, USA)

Bacteria from uranium mining waste pile: interactions with U(VI) 262
P. Panak (Dresden, Germany), B.C. Hard, K. Pietzsch (Leipzig, Germany), S. Kutschke, K. Röske, S. Selenska-Pobell, G. Bernhard, H. Nitsche (Dresden, Germany)

The investigation of the neptunium complexes formed upon interaction of high level waste glass and Boom Clay media 267
V. Pirlet, P. Van Iseghem, A. Dierckx (Mol, Belgium), J.-F. Desreux (Sart-Tilman, Belgium)

Solubility measurements and sorption studies of thorium in cement pore water 272
B. Wierczinski (Göteborg, Sweden), S. Helfer, M. Ochs (Zürich-Schlieren, Switzerland), G. Skarnemark (Göteborg, Sweden)

Heavy Element Research

Chemistry gains a new element: Z=106 277
H.W. Gäggeler (Berne and Villigen, Switzerland)

Recent progress in theoretical investigations of the electronic structure of the transactinides 283
V. Pershina (Darmstadt, Germany), T. Bastug (Kyoto, Japan), B. Fricke (Kassel, Germany)

Evidence for relativistic effects in the chemistry of element 104 287
A. Türler (Villigen, Switzerland), G.V. Buklanov (Dubna, Russia), B. Eichler (Villigen, Switzerland), H.W. Gäggeler (Villigen and Berne, Switzerland), M. Grantz, S. Hübener (Dresden, Germany), D.T. Jost (Villigen, Switzerland), V.Ya. Lebedev (Dubna, Russia), D. Piguet (Villigen, Switzerland), S.N. Timokhin, A.B. Yakushev, I. Zvara (Dubna, Russia)

Extraction of the fluoride-, chloride- and bromide complexes of the elements Nb, Ta, Pa, and 105 into aliphatic amines 292
W. Paulus, J.V. Kratz, E. Strub, S. Zauner (Mainz, Germany), W. Brüchle, V. Pershina, M. Schädel, B. Schausten (Darmstadt, Germany), J.L. Adams, K.E. Gregorich, D.C. Hoffman, M.R. Lane, C. Laue, D.M. Lee, C.A. McGrath, D.K. Shaughnessy, D.A. Strellis, E.R. Sylwester (Berkeley, CA, USA)

Production of plutonium tracers in ^{237}Np nuclear reactions with ^{3}He-ions in the energy range from 26 to 60 MeV 296
J. Aaltonen (Helsinki, Finland), P. Dendooven (Jyväskylä, Finland), E.A. Gromova, V.A. Jakovlev (St. Petersburg, Russia), W.H. Trzaska (Jyväskylä, Finland)

Chromatographic studies of Rf (element 104) with tributylphosphate (TBP) 300
W. Brüchle, E. Jäger, V. Pershina, M. Schädel, B. Schausten (Darmstadt, Germany), R. Günther, J.V. Kratz, W. Paulus, A. Seibert, P. Thörle, S. Zauner (Mainz, Germany), D. Schumann (Dresden, Germany), B. Eichler, H. Gäggeler, D. Jost, A. Türler (Villigen, Switzerland)

Review of the SISAK system in transactinide research recent developments and future prospects 303
J.P. Omtvedt, J. Alstad (Oslo, Norway), K. Eberhardt (Mainz, Germany), K. Fure (Oslo, Norway), R. Malmbeck (Göteborg, Sweden), M. Mendel, A. Nähler (Mainz, Germany), G. Skarnemark (Göteborg, Sweden), N. Trautmann, N. Wiehl (Mainz, Germany), B. Wierczinski (Göteborg, Sweden)

Sorption behaviour of rutherfordium and thorium from HCl/Hf containing aqueous solution 307
D. Schumann, H. Nitsche (Dresden, Germany), St. Taut (Rossendorf, Germany), D.T. Jost, H.W. Gäggeler (Villigen, Switzerland), A.B. Yakushev, G.V. Buklanov, V.P. Domanov, D.T. Lien (Dubna, Russia), B. Kubica, R. Misiak, Z. Szeglowski (Krakow, Poland)

First aqueous chemistry with seaborgium (element 106) 312
M. Schädel (Darmstadt, Germany)

Thermochromatography of heavy actinides adsorption of No-259 on Ti, V, Nb, Ta and Mo 316
S. Taut, S. Hübener (Dresden, Germany), B. Eichler, A. Türler (Villigen, Switzerland), H.W. Gäggeler (Villigen and Berne, Switzerland), S.N. Timokhin, I. Zvara (Dubna, Russia)

Fission mechanisms in the carbon-induced fission of uranium 322
A. Yokoyama, H. Baba, N. Takahashi, T. Saito (Osaka, Japan)

Experimental verification of two deformation paths in the mass division process of actinides 327
Y.L. Zhao (Tokyo and Ibaraki, Japan), I. Nishinaka, Y. Nagame, K. Tsukada (Ibaraki, Japan), M. Tanikawa, K. Sueki, Y. Oura (Tokyo, Japan), S. Ichikawa, H. Ikezoe (Ibaraki, Japan), T. Ohtsuki (Sendai, Japan), H. Kudo (Niigata, Japan), H. Nakahara (Tokyo, Japan)

High-Pressure and High-Temperature Properties

Thermodynamic aspects of heavy fermion compounds under high pressure 331
T. Kagayama, G. Oomi (Kumamoto, Japan)

Correlated f-electron materials near a $T=0$ magnetic–nonmagnetic boundary: implications for δ-Pu 335
J.D. Thompson, M.F. Hundley, J.L. Sarrao (Los Alamos, NM, USA)

Local density approximation versus generalized gradient approximation: full charge density study of the atomic volume of the light actinides 339
L. Vitos (Lyngby, Denmark and Budapest, Hungary), J. Kollár (Budapest, Hungary), H.L. Skriver (Lyngby, Denmark)

High-pressure X-ray diffraction studies of Cm–Bk alloys: contribution to the actinide pressure-phase diagram 342
S. Heathman (Karlsruhe, Germany), R.G. Haire (Oak Ridge, TN, USA)

Actinide nitrides and nitride–halides in high-temperature systems 347
T. Ogawa, F. Kobayashi, T. Sato (Ibaraki-ken, Japan), R.G. Haire (Oak Ridge, TN, USA)

Structure and dynamic properties of molten uranium trichloride 355
Y. Okamoto, F. Kobayashi, T. Ogawa (Ibaraki-ken, Japan)

High-temperature transuranium organometallic species from laser ablation of actinide–polymer mixtures 359
J.K. Gibson (Oak Ridge, TN, USA)

Local structure of (U,Zr)Pd_3 by EXAFS 363
M. Akabori, T. Ogawa, A. Itoh, H. Motohashi, H. Shiwaku (Ibaraki-ken, Japan)

Calculated lattice relaxation in Pu-Ga alloys 367
J.D. Becker (Los Alamos, NM, USA), B.R. Cooper (Morgantown, WV, USA), J.M. Wills, L. Cox (Los Alamos, NM, USA)

Phase transformation in Pu–Ga alloys at low temperature and under pressure: limit stability of the δ phase 370
P. Deloffre, J.L. Truffier, A. Falanga (Is-sur-Tille, France)

Stability of UNCl in LiCl–KCl eutectic melt 374
F. Kobayashi, T. Ogawa, Y. Okamoto, M. Akabori (Ibaraki-ken, Japan)

Pressure effect on the magnetic and crystallographic structures in the $U(Ni_{1-x}Cu_x)_2Ge_2$ system 378
E. Caspi, H. Shaked, H. Pinto, M. Melamud (Beer-Sheva, Israel), Z. Hu, O. Chmaissem, S. Short, J.D. Jorgensen (Argonne, IL, USA)

Preparation and structure of quaternary mixed sulphide $Li(Lu,M)S_2$ (M=U,Ce) 382
N. Sato, H. Masuda, T. Fujino, K. Yamada, M. Wakeshima (Sendai, Japan)

Unusual variation of temperature factor of uranium dioxide at high temperature 386
H. Serizawa, K. Fukuda, T. Shiratori (Ibaraki-ken, Japan), T. Fujino, N. Sato, K. Yamada (Sendai, Japan)

Effect of cations on lattice constants of $(M_yU_{1-y})O_{2.00}$ (M=Pu, Th, La) at low doped cation concentrations 391
T. Tsuji, M. Iwashita (Nagoya, Japan), T. Yamashita, K. Ohuchi (Ibaraki, Japan)

Enthalpies of formation of UGa_2 and UGa_3 by calorimetry 395
B. Prabhakara Reddy, R. Babu, K. Nagarajan, P.R. Vasudeva Rao (Kalpakkam, India)

Electrical conductivity of uranium–neptunium mixed oxides 400
T. Yamashita, K. Ohuchi (Ibaraki, Japan), T. Tsuji, T. Kato, M. Ochida, M. Iwashita (Nagoya, Japan)

Thermal expansion of neptunium–plutonium mixed oxides 404
T. Yamashita, H. Muto (Ibaraki, Japan), T. Tsuji, Y. Nakamura (Nagoya, Japan)

Magnetism

Synchrotron radiation as a probe of actinide magnetism (T7-Al 'Magnetism') 408
D.B. McWhan (Upton, NY, USA)

Orbital and spin moments in uranium compounds from X-ray magnetic circular dichroism 414
P.D. de Réotier, J.-P. Sanchez, A. Yaouanc (Grenoble, France)

Electronic properties of UPdSn diluted by Lu 418
L. Havela, R.V. Dremov, A.V. Andreev, V. Sechovský, J. Šebek (Prague, Czech Republic), K. Prokeš (Prague, Czech Republic and Amsterdam, The Netherlands)

Specific heat measurements on $U_2(Ru_{1-x}Rh_x)_3Si_5$ ternary silicides 423
B. Chevalier (Pessac, France), L. Piraux, E. Grivei (Louvain-la-Neuve, Belgium), J. Etourneau (Pessac, France)

Pressure effects on magnetism in the uranium and neptunium monopnictides 426
D. Braithwaite, A. Demuer (Grenoble, France), I.N. Goncharenko (Gif-sur-Yvette, France and Moscow, Russia), V. Ichas (Karlsruhe, Germany), J.-M. Mignot (Gif-sur-Yvette, France), J. Rebizant, J.C. Spirlet (Karlsruhe, Germany), O. Vogt (Zürich, Switzerland), S. Zwirner (Karlsruhe, Germany)

Observation of two low-energy responses in UPd_2Al_3—interaction of magnetism and superconductivity? 433
N. Sato, N. Aso (Sendai, Japan), B. Roessli (Grenoble, France), G.H. Lander (Karlsruhe, Germany), T. Komatsubara, Y. Endoh, O. Sakai (Sendai, Japan)

Transport properties of $(U_{1-x}Np_x)Pd_2Al_3$ and $(U_{1-x}Pu_x)Pd_2Al_3$ intermetallic solid solutions 437
F. Wastin, E. Bednarczyk, J. Rebizant (Karlsruhe, Germany)

Magnetic structures of the ternary stannide $U_2(Ni_{0.70}Pd_{0.30})_2Sn$ 444
D. Laffargue (Pessac and Gif-sur-Yvette, France), F. Bourée (Gif-sur-Yvette, France), B. Chevalier (Pessac, France), T. Roisnel (Gif-sur-Yvette, France), S. Bordère (Pessac, France)

Magnetic structure of the ternary germanide $U_3Cu_4Ge_4$ 448
S. Pechev, B. Chevalier (Pessac, France), D. Laffargue (Pessac and Gif-sur-Yvette, France), B. Darriet, J. Etourneau (Pessac, France)

Crystal structure and magnetic susceptibility of uranium palladium sulfide bronze, $U_xPd_3S_4$ 452
T. Fujino, N. Sato, K. Yamada, H. Masuda, M. Wakeshima (Sendai, Japan)

UFe_2Zn_{20}: a new uranium intermetallic compound 456
A.P. Gonçalves, J.C. Waerenborgh (Sacavém, Portugal), A. Amaro, M. Godinho (Lisboa, Portugal), M. Almeida (Sacavém, Portugal)

Preparation of uranium-based magnetic amorphous alloys by RF sputtering 459
Y. Homma, Y. Takakuwa (Sendai, Japan), Y. Shiokawa (Ibaraki, Japan), K. Suzuki (Sendai, Japan)

Magnetic order in NpO_2 and UO_2 studied by muon spin rotation 463
W. Kopmann, F.J. Litterst, H.-H. Klauß, M. Hillberg, W. Wagener (Braunschweig, Germany), G.M. Kalvius, E. Schreier, F.J. Burghart (Garching, Germany), J. Rebizant, G.H. Lander (Karlsruhe, Germany)

Magneto-optical investigations of URhAl intermetallic compound 467
M. Kučera, P. Beránková, M. Matyáš, I. Tichý (Prague, Czech Republic), A.A. Menovsky (Amsterdam, The Netherlands)

Magnetic properties of the $U_{1-x}Gd_xNi_2$ system 471
P. Lucaci, E. Burzo, I. Lupsa (Cluj-Napoca, Romania)

Incommensurate and ferrimagnetic phases in $U(Ni,M)_2Si_2$ with minor M=Co or Cu 474
M. Kuznietz, H. Pinto, E. Caspi, M. Melamud (Beer-Sheva, Israel), G. André, F. Bourée (Gif-sur-Yvette, France)

Crystal structure of distorted fluorite-type SrU_2O_6 with pentavalent uranium 479
C. Miyake (Osaka, Japan), T. Fujino (Sendai, Japan)

Critical electron scattering in UGe_2 near the magnetic phase transition induced by pressure 482
G. Oomi, T. Kagayama (Kumamoto, Japan), Y. Ōnuki (Osaka, Japan)

Anisotropic resistivity and giant magnetoresistance in UNi_2Ge_2 and UNiGa from ab initio calculations 486
A.Ya. Perlov, P.M. Oppeneer, V.N. Antonov, A.N. Yaresko (Dresden, Germany), B.Yu. Yavorsky (Kiev, Ukraine)

Preparation and electrical resistivity of $(Pu_{1-x}U_x)Sb$ and $(Pu_{1-x}Y_x)Sb$ single crystals 490
J. Rebizant, F. Wastin, C. Rijkeboer, E. Bednarczyk, P. Lefèbvre (Karlsruhe, Germany)

Magnetic phases and magnetoelastic phenomena in UNiGa under pressure 495
F. Honda (Kumamoto, Japan), K. Prokeš (Prague, Czech Republic and Amsterdam, The Netherlands), M. Olšovec (Amsterdam, The Netherlands and Grenoble, France), F. Bourdarot, P. Burlet (Grenoble, France), T. Kagayama, G. Oomi (Kumamoto, Japan), L. Havela, V. Sechovský, A.V. Andreev (Prague, Czech Republic), E. Brück, F.R. de Boer, A.A. Menovsky (Amsterdam, The Netherlands), M. Mihálik (Amsterdam, The Netherlands and Košice, Slovakia)

Structure, magnetic and electrical properties of the $UMn_{4-x}Fe_xAl_8$ system 499
W. Suski, K. Wochowski, D. Badurski, A. Gilewski, T. Mydlarz (Wrocław, Poland)

Ac-susceptibility and neutron diffraction studies of the $URu_{1-x}Pd_xGa$ system 503
V.H. Tran (Wroclaw, Poland), G. André, F. Bourée (Gif-Sur-Yvette, France), R. Troć (Wroclaw, Poland), H. Noël (Rennes, France)

High temperature susceptibilities of actinide monopnictides and monochalcogenides 508
O. Vogt, K. Mattenberger (Zürich, Switzerland), J. Löhle (Villigen, Switzerland), J. Rebizant (Karlsruhe, Germany)

On the characterization of $(U,Np)_xT_yX_z$ intermetallic solid solutions 513
F. Wastin, J. Rebizant, A.D. Stalios, C.T. Walker (Karlsruhe, Germany)

Nuclear Technology and Applications/Thermodynamic Properties

Nuclear fuels for actinides burning CAPRA and SPIN programmes status 517
A. Languille, P. Millet, J. Rouault, S. Pillon (Saint Paul-lez-Durance, France)

Actinide transmutation properties of thermal and fast fission reactors including multiple recycling 522
H.W. Wiese (Karlsruhe, Germany)

Concept of hydride fuel target subassemblies in a fast reactor core for effective transmutation of MA 530
M. Yamawaki, H. Suwarno (Tokyo, Japan), T. Yamamoto, T. Sanda, K. Fujimura, K. Kawashima, K. Konashi (Ibaraki-ken, Japan)

The applications of formo- and aceto-hydroxamic acids in nuclear fuel reprocessing 534
R.J. Taylor, I. May, A.L. Wallwork, I.S. Denniss (Sellafield, UK), N.J. Hill (Risley, UK), B.Ya. Galkin, B.Ya. Zilberman, Yu.S. Fedorov (St. Petersburg, Russia)

Separation of actinides and fission products in high-level liquid wastes by the improved TRUEX process 538
M. Ozawa, Y. Koma, K. Nomura, Y. Tanaka (Ibaraki-ken, Japan)

Inert matrices for the transmutation of actinides: fabrication, thermal properties and radiation stability of ceramic materials . . 544
M. Burghartz, Hj. Matzke, C. Léger (Karlsruhe, Germany), G. Vambenepe, M. Rome (Villeurbanne, France)

Mechanical and thermal properties of uranium intermetallic compounds 549
S. Yamanaka, K. Yamada, T. Tsuzuki, T. Iguchi, M. Katsura (Osaka, Japan), Y. Hoshino, W. Saiki (Ibaraki, Japan)

Actinide transmutation: new investigation on some actinide compounds 557
M. Beauvy, T. Duverneix, C. Berlanga, R. Mazoyer, C. Duriez (Saint-Paul-lez-Durance, France)

Development of transuranium elements recovery from high-level radioactive liquid waste 563
R. Fujita, Y. Akai (Kawasaki, Japan)

XPS and XRD studies of corrosion of uranium nitride by water 568
S. Sunder (Chalk River, Canada), N.H. Miller (Pinawa, Canada)

Influence of low temperature air oxidation on the dissolution behaviour of UO_2 and MOX spent fuel 573
J.A. Serrano (Madrid, Spain), J.P. Glatz, E.H. Toscano, D. Papaioannou (Karlsruhe, Germany), J. Barrero (Madrid, Spain), M. Coquerelle (Karlsruhe, Germany)

Thermophysical and thermodynamic properties of actinide mononitrides and their solid solutions 577
Y. Suzuki, Y. Arai (Ibaraki-ken, Japan)

The vibrational frequencies, molecular geometry and thermodynamic properties of the actinide tetrahalides 583
R.J.M. Konings (Petten, The Netherlands), D.L. Hildenbrand (Menlo Park, CA, USA)

Thermodynamics of reductive extraction of actinides and lanthanides from molten chloride salt into liquid metal 587
H. Moriyama, H. Yamana, S. Nishikawa, S. Shibata (Osaka, Japan), N. Wakayama, Y. Miyashita, K. Moritani (Kyoto, Japan), T. Mitsugashira (Sendai, Japan)

Measurement of standard potentials of actinides (U,Np,Pu,Am) in LiCl–KCl eutectic salt and separation of actinides from rare earths by electrorefining 592
Y. Sakamura, T. Hijikata, K. Kinoshita, T. Inoue (Tokyo, Japan), T.S. Storvick, C.L. Krueger (Columbia, MO, USA), J.J. Roy, D.L. Grimmett, S.P. Fusselman, R.L. Gay (Canoga Park, CA, USA)

Interdiffusion in the U–Zr system at δ-phase compositions 597
M. Akabori, A. Itoh, T. Ogawa (Ibaraki-ken, Japan), T. Ogata (Tokyo, Japan)

Thermal conductivity of actinide mononitride solid solutions 602
Y. Arai, K. Nakajima, Y. Suzuki (Ibaraki-ken, Japan)

The EFTTRA experiment on irradiation of Am targets 606
J-F. Babelot (Karlsruhe, Germany), R. Conrad, R.J.M. Konings (Petten, The Netherlands), G. Mühling (Karlsruhe, Germany), M. Salvatores (Saint-Paul-lez-Durance, France), G. Vambenepe (Villeurbanne, France)

Phase characterisation of simulated high burn-up UO_2 fuel 610
J. Cobos, D. Papaioannou, J. Spino, M. Coquerelle (Karlsruhe, Germany)

Fabrication of transmutation and incineration targets by infiltration of porous pellets by radioactive solutions 616
A. Fernandez, K. Richter, J. Somers (Karlsruhe, Germany)

Investigation of thorium(IV) separation from cerium(IV) on a silica gel column coated with tri-octylphosphine oxide 620
Ü. Hiçsönmez, M. Eral (İzmir, Turkey)

Thermodynamic study of the vaporization of Cs_2UO_4 by high temperature mass spectrometry 625
J. Huang, M. Yamawaki, K. Yamaguchi, F. Ono, M. Yasumoto (Tokyo, Japan), H. Sakurai, J. Sugimoto (Ibaraki-ken, Japan)

Hydrogen absorption properties of U_6Mn and U_6Ni 629
H. Ito (Tokyo, Japan), K. Yamaguchi, T. Yamamoto (Ibaraki, Japan), M. Yamawaki (Tokyo, Japan)

The influence of americium oxide inclusions on the thermal conductivity of inert-matrix fuel 632
K. Bakker, R.J.M. Konings (Petten, The Netherlands)

Thermodynamic assessment of the Fe–U, U–Zr and Fe–U–Zr systems 636
M. Kurata, T. Ogata, K. Nakamura (Tokyo, Japan), T. Ogawa (Ibaraki-ken, Japan)

Phase equilibria in the ternary URu_3–URh_3–UPd_3 system 641
K. Kurosaki, M. Uno (Osaka, Japan)

Modeling and estimating the distribution of trivalent actinides and lanthanides between molten fluorides and liquid metal . . . 645
F. Lemort, R. Piccinato, R. Boen (Bagnols-sur-Cèze, France), M. Allibert (Saint Martin d'Hères, France)

The formation of hydrophilic Np(IV) complexes and their potential application in nuclear fuel reprocessing 650
I. May, R.J. Taylor, G. Brown (Sellafield, UK)

Thermodynamics of nitrogen-rich uranium sesquinitride formation by reaction of uranium monocarbide with ammonia 654
K. Nishimaki (Osaka, Japan), M. Hirota (Hyogo, Japan), T. Nakagawa, T.A. Yamamoto (Osaka, Japan), M. Miyake (Fukui, Japan), M. Katsura (Osaka, Japan)

Thermodynamic study on α-U_2N_{3+x} using N-rich starting material ($x \geq 0.6$) 658
T. Nakagawa, K. Nishimaki, T. Urabe, M. Katsura (Osaka, Japan)

The formation of uranium sesquinitride by reactions of U or UH_3 with a N_2–H_2 mixture 662
M. Katsura, K. Nishimaki, T. Nakagawa, K. Takahashi (Osaka, Japan)

Vaporization behavior of (Np,Pu)N 666
K. Nakajima, Y. Arai, Y. Suzuki (Ibaraki-ken, Japan)

Irradiation behavior of microspheres of U-Zr alloys 670
T. Ogawa (Ibaraki-ken, Japan), T. Ogata (Tokyo, Japan), A. Itoh, M. Akabori, H. Miyanishi, H. Sekino, M. Nishi, A. Ishikawa (Ibaraki-ken, Japan)

Enthalpy of solution of californium oxychloride; calculation of the standard enthalpy of formation of CfOCl 676
J.B. Burns (Knoxville, TN, USA), R.G. Haire (Oak Ridge, TN, USA), J.R. Peterson (Knoxville and Oak Ridge, TN, USA)

Electrochemical preparation of high purity uranium and neptunium metals from aqueous solutions 680
K. Hasegawa, Y. Shiokawa (Ibaraki, Japan), M. Akabori (Ibaraki-ken, Japan), Y. Suzuki (Ibaraki, Japan), K. Suzuki (Sendai, Japan)

Electrochemical behavior of actinide ions in LiCl–KCl eutectic melts 685
O. Shirai, T. Iwai, Y. Suzuki (Ibaraki-ken, Japan), Y. Sakamura, H. Tanaka (Tokyo, Japan)

Neptunium concentration profiles in the Purex process 689
M. Takanashi, S. Homma, J. Koga, S. Matsumoto (Urawa, Japan)

Separation of actinides from simulated spent fuel solutions by an advanced ion-exchange process 693
Y.-Z. Wei, M. Yamaguchi, M. Kumagai, Y. Takashima (Kashiwa, Japan), T. Hoshikawa, F. Kawamura (Hitachi, Japan)

Mechanical properties of $(U,Ce)O_2$ 697
K. Yamada, S. Yamanaka, M. Katsura (Osaka, Japan)

Preparation, analysis and irradiation of hydrided U–Th–Zr alloy samples for a new fuel 702
T. Yamamoto (Sendai, Japan), H. Suwarno, F. Ono (Ibaraki, Japan), H. Kayano (Sendai, Japan), M. Yamawaki (Ibaraki, Japan)

Equilibrium distributions of trivalent actinides and lanthanides in the liquid–liquid extraction system of TBP and molten $Ca(NO_3)_2 \cdot 4H_2O$ 707
H. Yamana (Osaka, Japan), T. Kaibuki, Y. Miyashita (Kyoto, Japan), S. Shibata, H. Moriyama (Osaka, Japan)

Solution Chemistry, Including Thermodynamic Aspect

Kinetics of actinide complexation reactions 712
K.L. Nash, J.C. Sullivan (Argonne, IL, USA)

Comparative study on the hydration states of Cm(III) and Eu(III) in solution and in cation exchange resin 719
T. Kimura, Y. Kato, H. Takeishi (Ibaraki, Japan), G.R. Choppin (Tallahassee, FL, USA)

Recent achievements of actinide radiation chemistry 723
M.V. Vladimirova (Moscow, Russia)

Spectroscopic evaluation of thermodynamics of trivalent actinides in brines 728
Th. Fanghänel, J.-I. Kim (Karlsruhe, Germany)

Am (III)/Eu (III) separation at low pH using synergistic mixtures composed of carboxylic acids and neutral nitrogen polydendate ligands 738
P.Y. Cordier, C. Hill, P. Baron, C. Madic (Bagnols-sur-Cèze, France), M.J. Hudson (Whiteknights, UK), J.O. Liljenzin (Göteborg, Sweden)

Extraction of americium and lanthanides by dialkyldithiophosphinic acid and f-f absorption spectra of the extraction complexes 742
Z. Yongjun, X. Jingming, C. Jing, C. Yingzhong (Beijing, P.R. China)

A complexation study of Cm(III) and Tb(III) with chelating aromatic ligands by time-resolved laser fluorescence spectroscopy 746
R. Klenze (Karlsruhe, Germany), P. Panak (Dresden, Germany), J.I. Kim (Karlsruhe, Germany)

Complexation of uranyl(VI) with polyoxometalates in aqueous solutions 751
A. Saito (Tokyo, Japan), G.R. Choppin (Tallahassee, FL, USA)

Deactivation and luminescence lifetimes of excited uranyl ion and its fluoro complexes 756
Z. Fazekas, T. Yamamura, H. Tomiyasu (Tokyo, Japan)

New method for actinide and rare-earth element recovery by diphenyl[dibutylcarbamoylmethyl]phosphine oxide from nitric acid solutions 760
Y.M. Kulyako, D.A. Malikov, M.K. Chmutova, M.N. Litvina, B.F. Myasoedov (Moscow, Russia)

Interaction of uranium(VI) with silicic acid in aqueous solutions studied by time-resolved laser-induced fluorescence spectroscopy (TRLFS) 765
H. Moll, G. Geipel, V. Brendler, G. Bernhard, H. Nitsche (Dresden, Germany)

Distribution of uranium on zeolite X and investigation of thermodynamic parameters for this system 769
S. Akyıl, M.A.A. Aslani, Ş.(Ö.) Aytaş (Bornova-İzmir, Turkey)

Complexation of NpO_2^+ by aromatic polycarboxylates 774
G.R. Choppin (Tallahassee, FL, USA), B.E. Stout, M. Pagès (Paris, France)

Solvent extraction of U(VI) from media containing competing anions and metal cations by a calix[6]arene bearing carboxy groups 778
C. Dinse, N. Baglan, C. Cossonnet (Fontenay-aux-Roses, France), J.F. Le Du (Orsay, France), Z. Asfari, J. Vicens (Strasbourg, France)

Thermodynamics of actinide redox reactions in potassium phosphotungstate solutions 782
E.A. Erine, A.A. Baranov, A.Yu. Volkov, V.M. Chistyakov, G.A. Timofeev (Ulyanovsk, Russia)

Chloride and pseudohalide complexes of thorium in anhydrous solvents 786
V.A. Kalibabchuk, G.N. Zaitseva (Kiev, Ukraine)

Steric effect of β-diketone in synergistic extraction of actinide(III) and lanthanide(III) with β-diketone +18-crown-6 ether/1,2-dichloroethane 790
Y. Meguro, Y. Kitatsuji, T. Kimura, Z. Yoshida (Ibaraki, Japan)

Isotope and temperature effects on photochemical reactions of uranyl ion in H_2O–D_2O mixtures 794
R. Nagaishi, T. Kimura, J. Inagawa, Y. Kato (Ibaraki-ken, Japan)

Solvent extraction study of actinide elements by *N,N'*-dimethyl-*N,N'*-dihexyl-3-oxapentanediamide and thenoyltrifluoroacetone . 799
Y. Sasaki, T. Adachi (Ibaraki, Japan), G.R. Choppin (Tallahassee, FL, USA)

The interaction of Np(V), Pu(VI) and Tc(VII) with metal in alkaline solutions 803
V.I. Silin, A.V. Kareta (Moscow, Russia)

Complexation behavior of Pu(IV) and Pu(VI) with urea in nitric acid solution 809
S.I. Sinkov, E.I. Bozhenko (Moscow, Russia)

Behavior of plutonium(V) in alkaline media 813
N.A. Budantseva, I.G. Tananaev, A.M. Fedoseev (Moscow, Russia), C.H. Delegard (Richland, WA, USA)

The oxidation of neptunium(IV) by nitric acid in 100% TBP and diluted TBP/*n*-dodecane solutions 817
R.J. Taylor (Sellafield, UK), V.S. Koltunov, O.A. Savilova, G.I. Zhuravleva (Moscow, Russia), I.S. Denniss, A.L. Wallwork (Sellafield, UK)

Criteria for compilation of a site-specific thermodynamic database for geochemical speciation calculations 821
M. Chandratillake, D.P. Trivedi, M.G. Randall, P.N. Humphreys (Risley, UK), E.J. Kelly (Sellafield, UK)

Spectroscopy

ARPES in strongly correlated 4f and 5f systems: Comparison to the Periodic Anderson Model (PAM) 826
A.J. Arko, J.J. Joyce, L.E. Cox, L. Morales, J. Sarrao, J.L. Smith (Los Alamos, NM, USA), Z. Fisk (Tallahassee, FL, USA), A. Menovsky (Amsterdam, The Netherlands), A. Tahvildar-Zadeh, M. Jarrell (Cincinnati, OH, USA)

Optical and magneto-optical spectroscopy of uranium and plutonium compounds: recent theoretical progress 831
P.M. Oppeneer, A.Ya. Perlov, V.N. Antonov, A.N. Yaresko, T. Kraft (Dresden, Germany), M.S.S. Brooks (Karlsruhe, Germany)

Determination of the first ionization potential of nine actinide elements by resonance ionization mass spectroscopy (RIMS) . . 837
N. Erdmann, M. Nunnemann, K. Eberhardt, G. Herrmann, G. Huber, S. Köhler, J.V. Kratz, G. Passler (Mainz, Germany), J.R. Peterson (Knoxville, TN, USA), N. Trautmann, A. Waldek (Mainz, Germany)

Thin layers in actinide research 841
T. Gouder (Karlsruhe, Germany)

Synthesis and characterization of actinide-exchanged Preyssler heteropolyanions $[AnP_5W_{30}O_{110}]^{n-}$ (An≡Th, Am, Cm) 846
M.R. Antonio, C.W. Williams, L. Soderholm (Argonne, IL, USA)

Spectroscopy and dynamics of 5f states of Es^{3+} in LaF_3 850
J.V. Beitz, C.W. Williams, G.K. Liu (Argonne, IL, USA)

Emission profile of Bk(III) in a silicate matrix: anomalous dependence on excitation power 854
Z. Assefa, R.G. Haire (Oak Ridge, TN, USA), N.A. Stump (Winston-Salem, NC, USA)

Electronic structure of f^1 actinide complexes. Part 3 of [1]. Quasi-relativistic density functional calculations of the optical transition energies of PaX_6^{2-} (X=F, Cl, Br, I) 859
N. Kaltsoyannis (London, England)

Synthesis and characterization of strontium pentachlorouranate(III) and dibarium heptachlorouranate(III) 863
M. Karbowiak, J. Drożdżyński (Wrocław, Poland)

Luminescence study on determination of the inner-sphere hydration number of Am(III) and Nd(III) 867
T. Kimura, Y. Kato (Ibaraki, Japan)

Effects of self-radiation damage on electronic properties of $^{244}Cm^{3+}$ in an orthophosphate crystal of YPO_4 872
G.K. Liu, S.T. Li, J.V. Beitz (Argonne, IL, USA), M. Abraham (Oak Ridge, TN, USA)

Determination of the first ionization potential of einsteinium by resonance ionization mass spectroscopy (RIMS) 876
J.R. Peterson (Knoxville, TN, USA), N. Erdmann, M. Nunnemann, K. Eberhardt, G. Huber, J.V. Kratz, G. Passler, O. Stetzer, P. Thörle, N. Trautmann, A. Waldek (Mainz, Germany)

PuGa alloys: an X-ray absorption study 879
N. Richard, Ph. Faure, P. Rofidal, J.L. Truffier, D. Bazin (Is-sur-Tille, France)

Surface contamination of single-crystal PuSb 882
S.R. Wasserman, L. Soderholm (Argonne, IL, USA), J. Rebizant, G.H. Lander (Karlsruhe, Germany)

Author Index of Volumes 271–273 887

Subject Index of Volumes 271–273 891

CONTENTS DIRECT

ContentsDirect, the ***free*** e-mail alerting service, delivers the table of contents for this journal directly to your PC, prior to publication. The quickest way to register for **ContentsDirect** is via the Internet at: **http://www.elsevier.nl/locate/ContentsDirect**. If you don't have access to the Internet you can register for this service by sending an e-mail message to cdsubs@elsevier.co.uk – specifying the title of the publication you wish to register for.